# Real-World Utilization of Glucagon-Like Peptide-1 Receptor Agonists in Overweight and Obese US Adults



**RWD112** 

## Introduction

- Olucagon-like peptide-1 receptor agonists (GLP-1 RAs) are a cornerstone of type 2 diabetes and weight-loss treatments<sup>1,2</sup>
- Description Approvals of some GLP-1 RAs for weight-loss treatment have increased the popularity of these drugs, resulting in possible off-label usage
- Ourrently, multiple products utilizing GLP-1 RAs are approved by the US Food and Drug Administration for weight management concurrent with a reduced-calorie diet and increased physical activity<sup>3-5</sup>

\*Dave Iwanyckyj, BA1; Rohan Vashi, PharmD, MS1; Pablo Racana, BS1; Melanie Jardim, PhD1

GLP-1 RA: semaglutide (WEGOVY®)

<sup>1</sup>Amplity, Inc, Langhorne, PA, USA

- GLP-1 RA: liraglutide (SAXENDA®)
- Dual glucose-dependent insulinotropic peptide and GLP-1 RA tirzepatide (ZEPBOUND®)
- In the United States, these agents are indicated for patients with a body mass index (BMI) of
- ≥30 kg/m² OR
- 27 kg/m<sup>2</sup> to <30 kg/m<sup>2</sup> (overweight) and at least 1 weight-related comorbid condition
- Deight-related comorbid condition: hypertension, dyslipidemia, type 2 diabetes mellitus, obstructive sleep apnea, or cardiovascular disease
- Dayer rejection rates of GLP-1 RAs are high (62.4%), primarily attributed to prior authorization requirements and formulary exclusion<sup>6</sup>
- This research aims to characterize real-world usage of GLP-1 RAs among overweight and obese US adults

## Methods

- Descriptions of US prescriber-patient and platform built from HIPAA-compliant transcriptions of US prescriber-patient visits. Using AI and natural language processing (NLP), it extracts, visualizes, and summarizes treatment discussions and clinical decisions. Covering inpatient and outpatient care across 70+ specialties since 2017, AnswerY was known as *Amplity Insights™* prior to January 2025
- NLP was used to search and analyze the AnswerY database and platform for records from 12,540 providers for patients who had received a GLP-1 RA as a class or specific agent from January 1, 2017, to October 30, 2024
- Datient demographics and comorbidities were aggregated, and drug-utilization data from 193,193 records were summarized for overweight or obese patients with and without diabetes, and a BMI <25, 25-26, 27-29, and ≥30

# Conclusion

- In this real-world US cohort, GLP-1 RA-containing medications are used heavily in overweight and obese populations. However, usage of GLP-1 RAs among overweight and obese patients are overwhelmingly used in patients with diabetes
- O Although the analysis focuses on overweight and obese patients, out of the 124,400 patients utilizing a GLP-1 RA, 54,201 (43.6%) were identified as being overweight or obese. The remaining 56.4% of patients were either not obese or phrases suggesting weight status or BMI may have been missing
- Liraglutide was the most utilized GLP-1 RA among overweight and obese patients with or without diabetes. Furthermore, utilization patterns for dulaglutide were different compared with liraglutide and semaglutide
- As dulaglutide is only approved to improve glycemic control as an adjunct to diet and exercise for patients with type 2 diabetes, usage of dulaglutide in patients without diabetes suggests potential off-label usage<sup>7</sup>
- In combination with the popularity of GLP-1 RAs, potential off-label usage may lead to drug shortages for patients utilizing GLP-1 RAs for type 2 diabetes treatment Drug shortages for GLP-1 RAs such as the one seen from 2022-2024 may lead to pharmacies temporarily compounding GLP-1 RAs,
- which may have additional patient safety and efficacy concerns not associated with the original product • Increased manufacturing of brand and generic GLP-1 RAs, faster approvals of additional GLP-1 RAs and other agents in weight-loss treatment, and increased reimbursement for obesity treatment will provide a greater number of options for proper on-label usage of

#### \*PRESENTING AUTHOR

these agents



All authors are employees of Amplity, Inc.

To learn more about this study and further research by Amplity AnswerY™, please scan this code.

#### REFERENCES

- Hamasaki H. Exercise and glucagon-like peptide-1: does exercise potentiate the effect of treatment? World J Diabetes. 2018;9:138-140.
- Patoulias D, Popovic DS, Stoian AP, et al. Effect of semaglutide versus other glucagon-like peptide-1 receptor agonists on cardio-metabolic risk factors in patients with type 2 diabetes: a systematic review and meta-analysis of head-to-head, phase 3, randomized controlled trials. J Diabetes Complications. 2023;37:108529 3. Novo Nordisk. WEGOVY (semaglutide) [prescribing information]. US Food and Drug Administration. Accessed April 1, 2025. https://www.accessdata.fda.gov/drugsatfda\_
- 1. Novo Nordisk. SAXENDA (liraglutide) [prescribing information]. US Food and Drug Administration. Accessed April 1, 2025. https://www.accessdata.fda.gov/drugsatfda\_docs/ label/2024/206321s019lbl.pdf
- 5. Eli Lilly. ZEPBOUND (tirzepatide) [prescribing information]. US Food and Drug Administration. Accessed April 1, 2025. https://www.accessdata.fda.gov/drugsatfda\_docs/
- label/2025/217806Orig1s020lbl.pdf 6. Son M, Edwards A, Burns B. Yearly trends in payer coverage rates for GLP-1 receptor antagonists in weight loss. Presented at: AMCP 2025. March 31-April 3, 2025; Houston, TX.
- 7. Eli Lilly. TRULICITY (dulaglutide) [prescribing information]. US Food and Drug Administration. Accessed April 1, 2025. https://www.accessdata.fda.gov/drugsatfda\_docs/ label/2024/125469s061s062lbl.pdf


## Patients utilizing GLP-1 RAs mostly have diabetes, and 43.6% are overweight or obese

3.3%

# **Results: Patient Demographics**

- AnswerY identified 124,400 patients utilizing a GLP-1 RA from 19,627 providers from January 1, 2017, to October 30, 2024
- Patients utilizing GLP-1 RAs were older (age, mean ± standard deviation [SD]: 61 ± 12.8 years), and of patients with a known race (46.8%), 11.9% of patients were African-American, 0.4% were Asian, 4.7% were Hispanic, and 82.9% were White
- diabetes. Most patients with a recorded BMI had a BMI ≥30 (68.3%)
- Figure 1 shows market understanding of GLP-1 RA usage, and Table 1 shows patient demographics stratified by BMI within the identified cohort utilizing the AnswerY real-world database and platform


#### Figure 1: Summary of GLP-1 RA Utilizing Patient Demographics (n=124,400)





BMI, body mass index; GLP-1 RA, glucagon-like peptide-1 receptor agonist.


GLP-1 RA, glucagon-like peptide-1 receptor agonist.

89.9%










# Table 1: GLP-1 RA Utilizing Patient Demographics Stratified by BMI BMI <25 BMI 25-26 BMI 27-29 BMI ≥30 Patients With Record, n (%) 139 (65.3) 310 (66.1) 772 (64.3) 2,547 (65.4) Mean (SD) 63.2 (13.4) 61.8 (12.8) 61.0 (12.8) 56.6 (13.4) Patients With Record, n (%) 213 (100) 468 (99.8) 1,122 (99.9) 3,887 (100) Male, n (%) 79 (37.1) 198 (42.3) 484 (43.1) 1,650 (42.4) Female, n (%) 134 (62.9) 270 (57.7) 638 (56.9) 2,237 (57.6) Patients With Record. n (%) 79 (37.1) 166 (35.4) 381 (33.9) 1.302 (33.4) African-American, n (%) 12 (15.2) 27 (16.3) 11 (6.6) 30 (7.9) 68 (5.2) White, n (%) 64 (81.0) 128 (77.1) 286 (75.1) 1,059 (81.3) BMI, body mass index; GLP-1 RA, glucagon-like peptide-1 receptor agonist.

## Liraglutide is the most utilized GLP-1 RA among overweight and obese patients with and without diabetes, regardless of BMI

## **Results: GLP-1 RA Utilization**


- Figure 2 shows the most utilized GLP-1 RAs among patients with and without diabetes within the cohort
- Liraglutide was the most utilized GLP-1 RA among patients with and without diabetes, followed by dulaglutide and semaglutide
- Liraglutide (49.5% vs 44.5%) and semaglutide (35.6% vs 15.6%) both have higher utilization in patients without diabetes, whereas dulaglutide (15.5% vs 34.7%) shows greater utilization in patients with diabetes

#### Figure 2: GLP-1 RA Agent Usage Among Patients With and Without Diabetes



- Figure 3 shows the most utilized GLP-1 RAs among patients with a recorded BMI, stratified by BMI
- Utilization of GLP-1 RAs among BMI-stratified cohorts was similar to patients with and without diabetes, with liraglutide having higher overall utilization rates than other agents
- Among those that utilized dulaglutide, patients with a BMI of <25 utilized dulaglutide more than patients with higher BMIs</p> (BMI <25 vs BMI ≥30: 39.0% vs 32.5%)

#### Figure 3: GLP-1 RA Agent Usage Stratified by BMI




#### Patients with diabetes utilizing GLP-1 RAs show higher rates of most comorbidities compared with patients without diabetes

## **Results: Comorbidities**

- Ocomorbidities were present in 82.1% and 64.9% of patients with and without diabetes, respectively. Figure 4 shows the most common comorbidities
- Decompared with patients with diabetes, patients without diabetes experience overall lower rates of all comorbidities except for mental health disorders (36.5% vs 47.9%)

#### Figure 4: Prevalence of Comorbidities Among Patients With and Without Diabetes



Some patients utilizing GLP-1 RAs utilize SGLT2i's. Among those patients, regardless of having diabetes or a patient's BMI, empagliflozin was the most common SGLT2i utilized

Prevalence of Comorbidity (%)

## Results: SGLT2i Usage

- Oscilium-glucose cotransporter-2 inhibitors (SGLT2i's) were used among 16.5% of all patients in the cohort. **Table 2** shows the usage of the most common SGLT2i's among all overweight and obese patients with and without diabetes, and stratified by BMI
- Among all patients and the entire cohort, empagliflozin was the most utilized concomitant medication

#### Table 2: SGLT2i Usage Among Patients Utilizing GLP-1 RAs

	All Patients	Overweight and Obese With Diabetes	Overweight and Obese Without Diabetes	BMI <25	BMI 25-26	BMI 27-29	BMI ≥30
All Patients (%)	100	100	100	100	100	100	100
Any SGLT2i (%)	16.5	17.4	3.6	20.7	21.5	21.5	19.7
Canagliflozin (%)	25.3	26.2	23.8	18.2	23.6	23.8	26.4
Dapagliflozin (%)	23.4	23.9	19	22.7	22.8	25.6	20.3
Empagliflozin (%)	57.3	56.4	58.5	68.2	62.4	59.5	59.8

BMI, body mass index; GLP-1 RA, glucagon-like peptide-1 receptor agonists; SGLT2i, sodium-glucose cotransporter-2 inhibitor.

# Strength

Deliange number of overweight and obese patients found in the AnswerY real-world database and platform bolsters confidence in the robustness of the results

#### I Limitation

Description of the purchases, mail order, or weight-loss clinics, would not be captured in this analysis unless they provided information to their healthcare provider